

# Black Hole Radiation with Modified Dispersion Relation in Tunneling Paradigm: Free-fall Frame

Peng Wang,<sup>\*</sup> Haitang Yang,<sup>†</sup> and Shuxuan Ying<sup>‡</sup>

*Center for Theoretical Physics, College of Physical Science and Technology,*

*Sichuan University, Chengdu, 610064, PR China*

## Abstract

Due to the exponential high gravitational red shift near the event horizon of a black hole, it might appear that the Hawking radiation would be highly sensitive to some unknown high energy physics. To study effects of any unknown physics at the Planck scale on the Hawking radiation, the dispersive field theory models have been proposed, which are variations of Unruh's sonic black hole analogy. In this paper, we use the Hamilton-Jacobi method to investigate the dispersive field theory models. The preferred frame is the free-fall frame of the black hole. The dispersion relation adopted agrees with the relativistic one at low energy but is modified near the Planck mass  $m_p$ . The corrections to the Hawking temperature are calculated for massive and charged particles to  $\mathcal{O}(m_p^{-2})$  and neutral and massless particles with  $\lambda = 0$  to all orders. The Hawking temperature of radiation agrees with the standard one at the leading order. After the spectrum of radiation near the horizon is obtained, we use the brick wall model to compute the thermal entropy of a massless scalar field near the horizon of a 4D spherically symmetric black hole and a 2D one. Finally, the luminosity of a Schwarzschild black hole is calculated by using the geometric optics approximation.

---

<sup>\*</sup>Electronic address: pengw@scu.edu.cn

<sup>†</sup>Electronic address: hyanga@scu.edu.cn

<sup>‡</sup>Electronic address: 2013222020007@stu.scu.edu.cn

## Contents

<b>I. Introduction</b>	2
<b>II. Deformed Hamilton-Jacobi Method</b>	4
A. Deformed Hamilton-Jacobi Equation	4
B. Massive and Charged Particle to $\mathcal{O}(m_p^{-2})$	7
C. Massless and Neutral Particle to All Orders	9
D. Discussion	10
<b>III. Entropy in Brick Wall Model</b>	10
A. 4D Spherically Symmetric Black Hole	11
B. 2D Black Hole in Free-fall Scenario	14
C. 2D Black Hole in Static Scenario	15
D. Discussion	17
<b>IV. Black Hole Evaporation</b>	21
<b>V. Conclusion</b>	24
<b>Acknowledgments</b>	25
<b>References</b>	25

## I. INTRODUCTION

Soon after Stephen Hawking demonstrated that quantum effects could allow black holes to radiate a thermal flux of quantum particles[1], it was realized that there was the trans-Planckian problem with the calculation[2]. Hawking radiation appears to come from the modes with huge initial frequencies, well beyond the Planck mass  $m_p$ , which experience exponential high gravitational red-shifting near the horizon. So the Hawking radiation relies on the validity of quantum field theory in curved spacetime to arbitrary high energies. On the other hand, quantum field theory is considered more like an effective field theory of an underlying theory whose nature remains unknown. This observation poses the question

of whether any unknown physics at the Planck scale could strongly influence the Hawking radiation.

To study the trans-Planckian problem, a hydrodynamic analogue of a black hole radiation was considered[3]. Following the Unruh's work, there have been a lot of studies to understand the dispersive field theory models[3–13], which focused on studying the effect on the Hawking radiation due to modifications of the dispersion relations of matter fields at high energies. Similar to the original method for deriving the Hawking radiation, the energy fluxes for outgoing radiation were usually obtained by calculating the Bogoliubov transformations between the initial and final states of incoming and outgoing radiation. In most works, the Hawking effect could be recovered at leading order under some suitable assumptions, which have been briefly reviewed in [14, 15].

After the Hawking's original derivation, there have been some other methods proposed to understand the Hawking radiation. Recently, a semiclassical method of modeling Hawking radiation as a tunneling process has been developed and attracted a lot of attention. This method was first proposed by Kraus and Wilczek[16, 17], which is known as the null geodesic method. They employed the dynamical geometry approach to calculate the imaginary part of the action for the tunneling process of s-wave emission across the horizon and related it to the Hawking temperature. Later, the tunneling behaviors of particles were investigated using the Hamilton-Jacobi method[18–20]. In the Hamilton-Jacobi method, one ignores the self-gravitation of emitted particles and assumes that its action satisfies the relativistic Hamilton-Jacobi equation. The tunneling probability for the classically forbidden trajectory from inside to outside the horizon is obtained by using the Hamilton-Jacobi equation to calculate the imaginary part of the action for the tunneling process. Using the null geodesic method and Hamilton-Jacobi method, much fruit has been achieved[21–32]. Furthermore, the effects of quantum gravity on the Hawking radiation have been discussed in the Hamilton-Jacobi method. In fact, the minimal length deformed Hamilton-Jacobi equation for fermions in curved spacetime have been introduced and the modified Hawking temperatures have been derived[33–38]. These have motivated us to use the Hamilton-Jacobi method to study the dispersive field theory models[39]. In this paper, we focus on the dispersive models with the free-fall preferred frame, whereas those with the static preferred frame have been studied in [39].

The remainder of our paper is organized as follows. In section II, the deformed Hamilton-

Jacobi equations are derived for the dispersive models with the free-fall preferred frame. We then solve the deformed Hamilton-Jacobi equations to obtain tunneling rates for massive and charged particles to  $\mathcal{O}(m_p^{-2})$  and massless and neutral particles to all orders. The thermal entropy of a massless scalar field near the horizon is computed in section III using the brick wall model. In section IV, we calculate the luminosity of a Schwarzschild black hole with the mass  $M \gg m_p$ . Section V is devoted to our conclusion. Throughout the paper we take Geometrized units  $c = G = 1$ , where the Planck constant  $\hbar$  is square of the Planck mass  $m_p$ .

## II. DEFORMED HAMILTON-JACOBI METHOD

In this section, we first derive the deformed Hamilton-Jacobi equation incorporating the modified dispersion relation (MDR) assuming that the preferred reference frame is the free-fall frame. We then solve the deformed Hamilton-Jacobi equation for the imaginary part of  $I$  which gives the tunneling rate  $\Gamma$  across the event horizon. We consider two cases, a massive and charged particle to  $\mathcal{O}(m_p^{-2})$  and a neutral and massless particle with  $\lambda = 0$  to all orders.

### A. Deformed Hamilton-Jacobi Equation

To study the deformed Hamilton-Jacobi method incorporating the MDR for the Hawking radiation, one first needs to choose the form of MDR in flat spacetime (the local free-fall frame) and generalizes it to curved spacetime. To be as general as possible, we will work with the MDR for a particle with mass  $m$

$$E^2 = F^2(p) + m^2, \quad (1)$$

where we define

$$F(p) = p \sum_{n=0} C_n \frac{p^{2n}}{m_p^{2n}}, \quad (2)$$

$m_p$  is the Planck mass,  $C_0 = 1$ , and  $E$  and  $p$  are the energy and the norm of the momentum measured in some preferred reference frame, respectively. Note that the MDR (1) is rotational invariant in 4D spacetime. To generalize the MDR (1) to curved spacetime with the metric  $g_{\mu\nu}$ , we denote by  $u^\mu$  the unit vector field tangent to the observers' world lines,

which picks up a preferred frame. For a particle with the energy-momentum vector  $p_\mu$ , the energy  $E$  and the norm of the momentum  $p$  of the particle measured by these observers are

$$E = p_\mu u^\mu, \quad (3)$$

$$p^2 = E^2 - p_\mu p^\mu. \quad (4)$$

The curved spacetime generalization of the MDR (1) with a preferred frame described by  $u^\mu$  is obtained by plugging eqns. (3) and (4) into eqn. (1). To obtain the deformed Hamilton-Jacobi equation incorporating the MDR, one needs to relate the classical action  $I$  to  $p_\mu$ . In fact, it can be shown that, if  $I$  is a solution of the Hamilton-Jacobi equation, then the transformation equations give

$$p_\mu = -\partial_\mu I, \quad (5)$$

where  $-$  appears since  $p_\mu = (E, -\vec{p})$  in our metric signature. Replacing  $p_\mu$  with  $I$  via eqn. (5) and putting eqns. (3) and (4) into eqn. (1) gives the deformed Hamilton-Jacobi equation. In the appendix of [39], the deformed Hamilton-Jacobi equation is also derived in the language of the effective field theory for a scalar field and a fermion one. There we considered a scalar/fermion with the mass  $m$  and charge  $q$  in a static black hole with the presence of electromagnetic potential  $A_\mu$ . Neglecting self-interacting effective operators, we constructed the  $U(1)$  gauge invariant effective field theory incorporating the MDR. The deformed Klein-Gordon/Dirac equation was derived. The deformed Hamilton-Jacobi equation for scalars/fermions was then obtained using the WKB approximation. It was found there that the deformed scalar/fermionic Hamilton-Jacobi equation with respect to the preferred frame  $u^\mu$  in the black hole background spacetime can be written as

$$T^2 = F^2(X) + m^2, \quad (6)$$

where

$$T = -u^\mu (\partial_\mu I + qA_\mu), \quad X^2 = T^2 - (\partial_\mu I + qA_\mu)^2, \quad (7)$$

$A_\mu$  is the black hole's electromagnetic potential and  $q$  is the particle's charge.

As in [39], we here consider the black hole whose metric in the Schwarzschild-like coordinate is given by

$$ds^2 = f(r) dt^2 - \frac{dr^2}{f(r)} - C(r^2) h_{ab}(x) dx^a dx^b, \quad (8)$$

where  $f(r)$  has a simple zero at  $r = r_h$  with  $f'(r_h)$  being finite and nonzero. The vanishing of  $f(r)$  at point  $r = r_h$  indicates the presence of an event horizon. We also assumed that the black hole is asymptotically flat which gives  $f(r) \rightarrow 1$  as  $r \rightarrow \infty$ . However, a more suitable coordinate for describing a specific family of freely falling observers is the Painleve-Gullstrand (PG) coordinate[5, 6, 40]. The PG coordinate anchored to the freely falling observers along the radial direction takes the form of

$$ds^2 = dt_p^2 - [dr - v(r) dt_p]^2 - C(r^2) h_{ab}(x) dx^a dx^b, \quad (9)$$

where  $v(r)$  is the velocity of the free fall observer with respect to the rest observer and  $t_p$  measures proper time along them. The spacetime also has the event horizon at  $r_h$  satisfying  $v(r_h) = -1$ . We assume  $v < 0$ ,  $dv/dr > 0$  and  $v \rightarrow v_0 \leq 0$  as  $r \rightarrow \infty$ . Note that  $v < 0$  means the infalling observers. For simplicity we specialize to the particular family of observers with  $v_0 = 0$  who start at infinity with a zero initial velocity. Since the vector field  $u^\mu$  of the freely falling observers is tangent to the infalling world lines, one has for the infalling observers along the radial direction with  $v_0 = 0$  that

$$u^\mu = (1, v(r), \vec{0}) \quad (10)$$

in the PG coordinate and

$$u^\mu = \left( \frac{1}{f(r)}, \sqrt{1 - f(r)}, \vec{0} \right) \quad (11)$$

in the Schwarzschild-like coordinate. The fact that  $t_p$  is the proper time along the infalling world lines means that  $u_\mu$  is equal to the gradient of  $t_p$ ,

$$u_\mu = \partial_\mu t_p. \quad (12)$$

Using eqns. (11) and (12) gives

$$t_p = t + \int \frac{\sqrt{1 - f(r)}}{f(r)} dr. \quad (13)$$

Substituting eqn. (13) into the PG metric (9) and comparing it to the Schwarzschild-like coordinate (8), one finds

$$v(r) = -\sqrt{1 - f(r)}. \quad (14)$$

In the PG coordinate, one can use eqn. (10) to show that eqn. (7) becomes

$$\begin{aligned} T &= -(\partial_t I + qA_t) - v(r)(\partial_r I + qA_r), \\ X^2 &= (\partial_r I + qA_r)^2 + \frac{h^{ab}(x)(\partial_a I + qA_a)(\partial_b I + qA_b)}{C(r^2)}. \end{aligned} \quad (15)$$

Note that  $\partial_t$  and  $\partial_{t_p}$  are Killing vectors in the Schwarzschild-like coordinate and the PG coordinate, respectively. For the energy-momentum vector  $p_\mu$ , eqn. (13) gives us that its Killing energies associated with  $\partial_t$  and  $\partial_{t_p}$  are the same. Let  $\omega$  denote the Killing energies in the PG and Schwarzschild-like coordinates. Explicitly, one has  $\omega = \partial_t^\mu p_\mu = \partial_{t_p}^\mu p_\mu$ , which is a constant.

### B. Massive and Charged Particle to $\mathcal{O}(m_p^{-2})$

Since  $\omega = -\partial_{t_p} I$  is the conserved energy of the particle, we can separate  $t$  from other variables. Thus, we employ the following ansatz for the action  $I$

$$I = -\omega t + W(r) + \Theta(x). \quad (16)$$

The vector potential  $A_\mu$  is assumed to be given by

$$A_\mu = A_t(r) \delta_{\mu t}, \quad (17)$$

which is true for charged static black holes in most cases. Putting the ansatz (16) into eqn. (15), we have

$$\begin{aligned} T &= \tilde{\omega}(r) - v(r) p_r, \\ X^2 &= p_r^2 + \frac{h^{ab}(x) \partial_a \Theta(x) \partial_b \Theta(x)}{C(r^2)}, \end{aligned} \quad (18)$$

where  $\tilde{\omega}(r) = \omega - qA_t(r)$  and  $p_r = \partial_r W$ . The method of separation of variables gives the differential equation for  $\Theta(x)$

$$h^{ab}(x) \partial_a \Theta(x) \partial_b \Theta(x) = \lambda, \quad (19)$$

where  $\lambda$  is a constant and determined by  $h^{ab}(x)$ . Thus, one has

$$X^2 = p_r^2 + \frac{\lambda}{C(r^2)}, \quad (20)$$

and eqn. (6) becomes an ordinary differential equation for  $W(r)$ .

We consider a particle with mass  $m$  and charge  $q$ . Solving eqn.(6) for  $p_r$  gives

$$p_r^\pm = \frac{\pm \sqrt{\Lambda} + \sqrt{1-f}}{f} \tilde{\omega}(r) \mp \frac{C_1 \left[1 \pm \sqrt{(1-f)\Lambda}\right]^4 \tilde{\omega}^3(r)}{m_p^2 f^4 \sqrt{\Lambda}} + \mathcal{O}(m_p^{-4}), \quad (21)$$

where  $+/-$  denotes the outgoing/ingoing solutions and  $\Lambda = 1 - \frac{m^2 + \frac{\lambda}{C(r^2)}}{\tilde{\omega}^2(r)}$ . Here,  $p_r^+$  has a pole at  $r = r_h$ . To obtain the residue of  $p_r^+$  at  $r = r_h$ , one expands  $f(r)$  and  $C(r^2)$  at  $r = r_h$

$$\begin{aligned} f(r) &\sim 2\kappa(r - r_h) [1 + \eta\kappa(r - r_h) + \theta\kappa^2(r - r_h)^2 + \rho\kappa^3(r - r_h)^3], \\ C(r^2) &\sim C(r_h^2) [1 + c_1\kappa(r - r_h) + c_2\kappa^2(r - r_h)^2 + c_3\kappa^3(r - r_h)^3], \\ \tilde{\omega}(r) &\sim \tilde{\omega}(r_h) [1 + \omega_1\kappa(r - r_h) + \omega_2\kappa^2(r - r_h)^2 + \omega_3\kappa^3(r - r_h)^3] \end{aligned} \quad (22)$$

where  $\kappa = f'(r_h)/2$ . Using the residue theory for the semi circles, we get

$$\text{Im } W_+(r) = \frac{\tilde{\omega}(r_h)\pi}{\kappa} (1 + \Delta), \quad (23)$$

where we define

$$\begin{aligned} \Delta &= \frac{C_1}{2m_p^2} \left[ \frac{\delta_\lambda \lambda}{C(r_h^2)} + \delta_m m^2 + \delta_\omega \tilde{\omega}^2(r_h) \right] + \mathcal{O}(m_p^{-4}), \\ \delta_\lambda &= 3 + 2c_1^2 - 2c_2 + 12\eta + 12\eta^2 - 6\theta + c_1(6 + 6\eta - 2\omega_1) - 6\omega_1 - 6\eta\omega_1 + 2\omega_2, \\ \delta_m &= -1 + 12\eta^2 - 6\theta + \eta(4 - 6\omega_1) - 2\omega_1 + 2\omega_2, \\ \delta_\omega &= 40\eta^3 - 12\theta + 8\rho + \eta^2(24 - 60\omega_1) - 3\omega_1 + 24\theta\omega_1 + 12\omega_1^2 \\ &\quad - 2\omega_1^2 + 12\omega_2 - 12\omega_1\omega_2 + \eta(2 - 40\theta - 36\omega_1 + 24\omega_1^2 + 24\omega_2) - 6\omega_3. \end{aligned} \quad (24)$$

On the other hand, one can use eqns. (22) to expand  $p_r^-$  at  $r = r_h$ . It turns out that the residue of  $p_r^-$  at  $r = r_h$  is zero. Hence, we have  $\text{Im } W_-(r) = 0$ . As shown in [39], the probability of a particle tunneling from inside to outside the horizon is

$$P_{emit} \propto \exp \left[ -\frac{2}{\hbar} (\text{Im } W_+ - \text{Im } W_-) \right]. \quad (25)$$

There is a Boltzmann factor in  $P_{emit}$  with an effective temperature, which is

$$T_{eff} = \frac{T_0}{1 + \Delta}. \quad (26)$$

It is interesting to note that we have calculated  $\Delta$  in the static preferred frame in [39]. For emitted particles with mass  $m$  and charge  $q$ , we found

$$\Delta = -\frac{C_1}{2m_p^2} \left[ 2(3\omega_1 - \eta)\tilde{\omega}^2(r_h) + \frac{\lambda}{C(r_h^2)} + m^2 \right] + \mathcal{O}(m_p^{-4}). \quad (27)$$



### C. Massless and Neutral Particle to All Orders

We now work with a particle with  $m = 0$  and  $q = 0$ . To get all order result, we let  $\lambda = 0$ . Note that one has  $\lambda = 0$  for the solution in a 2D black hole or the s-wave solution in a 4D spherically symmetric black hole. In this case, we can use the following ansatz for the action  $I$

$$I = -\omega t + W(r). \quad (28)$$

Hence, the deformed Hamilton-Jacobi equation becomes

$$(\omega - vp_r)^2 = p_r^2 \left( \sum_{n=0}^{\infty} C_n \frac{p_r^{2n}}{m_p^{2n}} \right)^2, \quad (29)$$

where  $p_r = \partial_r W$ . We will prove by induction that the solutions to eqn. (29) take the form of

$$p_r^{\pm} = \frac{\pm\omega}{1 \pm v} \left[ 1 + \sum_{i=1}^{\infty} \frac{\omega^{2i}}{m_p^{2i}} \sum_{j=0}^{i-1} \frac{C_{i,j}}{(1 \pm v)^{3i-j}} \right], \quad (30)$$

where  $C_{i,j}$  is determined by  $C_n$ . In fact, it is easy to see that

$$p_r^{\pm} = \frac{\pm\omega}{1 \pm v} \left[ 1 - \frac{C_1 \omega^2}{m_p^2 (1 \pm v)^3} + \frac{\omega^4 [3C_1^2 - C_2 (1 \pm v)]}{m_p^4 (1 \pm v)^6} + \mathcal{O}\left(\frac{\omega^6}{m_p^6}\right) \right]. \quad (31)$$

Suppose for some integer  $N > 1$  such that

$$p_{r,N}^{\pm} = \frac{\pm\omega}{1 \pm v} \left[ 1 + \sum_{i=1}^N \frac{\omega^{2i}}{m_p^{2i}} \sum_{j=0}^{i-1} \frac{C_{i,j}}{(1 \pm v)^{3i-j}} \right] \quad (32)$$

satisfy eqn. (29) at  $\mathcal{O}\left(\frac{\omega^{2N}}{m_p^{2N}}\right)$ . Plugging  $p_r = p_{r,N}^{\pm} + \frac{\alpha^{\pm} \omega^{2N+2}}{m_p^{2N+2}} \frac{\omega}{1 \pm v}$  into eqn. (29), one finds that the terms of  $\mathcal{O}\left(\frac{\omega^{2N+2}}{m_p^{2N+2}}\right)$  gives

$$\alpha^{\pm} = \frac{P_{N-1}(1 \pm v)}{(1 \pm v)^{3(N+1)}}, \quad (33)$$

where  $P_{N-1}(x)$  is some polynomial of  $x$  with degree  $N-1$ . This completes the proof that eqn. (30) are solutions to eqn. (29) to all orders. We define the residue of  $\frac{1}{(1+v)^n}$  at  $r = r_h$  as

$$\text{Res}\left(\frac{1}{(1+v)^n}, r_h\right) = \frac{R_n}{\kappa}, \quad (34)$$

where  $R_1 = 1$ . Thus, one obtains

$$\text{Im } W^+ = \frac{\omega\pi}{\kappa} (1 + \Delta), \quad \text{Im } W^- = 0, \quad (35)$$

where we define

$$\Delta = \sum_{i=1}^{\infty} \frac{\delta_i \omega^{2i}}{m_p^{2i}} \text{ with } \delta_i = \left( \sum_{j=0}^{i-1} C_{i,j} R_{3i+1-j} \right). \quad (36)$$

Using eqn. (25) gives the effective temperature

$$T_{eff} = \frac{T_0}{1 + \Delta}. \quad (37)$$

## D. Discussion

When we use the residue theory for the semi circles to give eqns. (23) and (35), an assumption proposed in [39] is needed. The assumption requires that the singularity structure of  $\partial_r I$  except the order of the pole at  $r = r_h$  do not change after the MDR is introduced. It follows that  $\omega \lesssim m_p$ . A complete theory of quantum gravity might be needed to justify this assumption.

In most works of the dispersive models, much attention have been paid to the modifications of the asymptotic spectrum. Since the tunneling across the horizon takes place near the horizon, the near horizon spectrum of radiations is computed in the paper. The effects of scattering off the background need to be included if the asymptotic spectrum at infinity is considered. In most works, 2D spacetime has been considered. The higher order terms in the MDR violate conformal invariance of 2D spacetime, hence there is some scattering. Our calculations show that the spectrum of radiation near the horizon is close to a perfect thermal spectrum in the dispersive models. The thermal asymptotic spectrum has been recovered at leading order in previous studies. Thus, the energy fluxes of radiations are not significantly affected by the effects of scattering off the background. However, as noted in [39], the scattering effects might dramatically change the spectrum of radiations in the dispersive models with the static preferred frame.

## III. ENTROPY IN BRICK WALL MODEL

Bekenstein and Hawking showed that the entropy of a black hole is proportional to the area of the horizon[42–44]. Although all the evidences suggest that the Bekenstein-Hawking entropy is the thermodynamic entropy, the statistical origin of the black hole entropy has not yet been fully understood. It appears that an unavoidable candidate for the statistical

origin is the entropy of the thermal atmosphere of the black hole.

However, the entropy diverges when we attempt to calculate the entropy of the thermal atmosphere. There are two kinds of divergences. The first one is due to infinite volume of the system, which has to do with the contribution from the vacuum surrounding the system at large distances and is of little relevance here. The second one arises from the infinite volume of the deep throat region near the horizon. To regulate the divergences, t' Hooft [45] proposed the brick wall model for a scalar field  $\phi$ , where two brick wall cutoffs are introduced at some small distance  $r_\epsilon$  from the horizon and at a large distance  $L \gg r_h$ ,

$$\phi = 0 \quad \text{at } r = r_h + r_\epsilon \text{ and } r = L. \quad (38)$$

In the following, we will use the brick wall model to calculate the entropy of a scalar field for a 4D spherically symmetric black hole and a 2D one. For a 4D spherically symmetric black hole, the entropy will be calculated to  $O(m_p^{-2})$ . For the 2D black hole, we will obtain all order results in the cases with the static and free-fall preferred frames for comparison. For simplicity, we assume that the scalar field is massless and neutral.

### A. 4D Spherically Symmetric Black Hole

For a 4D spherically symmetric black hole with the Schwarzschild-like coordinate

$$ds^2 = f(r) dt^2 - \frac{dr^2}{f(r)} - C(r^2) (d\theta^2 + \sin^2 \theta d\phi^2), \quad (39)$$

we have shown that  $\lambda = (l + \frac{1}{2})^2 \hbar^2$  with the angular momentum  $l = 0, 1, \dots$  and the corresponding degeneracy is  $2l + 1$  [39]. Thus, the atmosphere entropy of a massless scalar field can be expressed in the form of

$$S = \int (2l + 1) dl \int d\omega \frac{dn(\omega, l)}{d\omega} s_{\omega, l}, \quad (40)$$

where  $\omega$  is the Killing energy associated with  $t$ ,  $l$  is the angular momentum,  $n(\omega, l)$  is the number of one-particle states not exceeding  $\omega$  with fixed value of angular momentum  $l$ , and  $s_{\omega, l}$  is the thermal entropy per mode. Taking the MDR corrections to both  $n(\omega, l)$  and the Hawking temperature into consideration, we used the brick wall model to calculate this entropy to all orders in [39], where the static preferred frame is used to import a MDR to the black hole background. By contrast, here the MDR corrected atmosphere entropy of the black hole is computed to  $\mathcal{O}(m_p^{-2})$  in the free-fall scenario.

For particles emitted in a wave mode labelled by energy  $\omega$  and  $l$ , we find that

(Probability for a black hole to emit a particle in this mode)

$$= \exp\left(-\frac{\omega}{T_{eff}}\right) \times (\text{Probability for a black hole to absorb a particle in the same mode}),$$

where  $T_{eff}$  is given by eqn. (26). The above relation for the usual dispersion relation was obtained by Hartle and Hawking [50] using semiclassical analysis. Neglecting back-reaction, detailed balance condition requires that the ratio of the probability of having  $N$  particles in a particular mode to the probability of having  $N - 1$  particles in the same mode is  $\exp\left(-\frac{\omega}{T_{eff}}\right)$ . One then follows the argument in [39] to get the average number  $n_{\omega,l}$  in the mode

$$n_{\omega,l} = n\left(\frac{\omega}{T_{eff}}\right), \quad (41)$$

where we define

$$n(x) = \frac{1}{\exp x - (-1)^\epsilon}. \quad (42)$$

Note that  $\epsilon = 0$  for bosons and  $\epsilon = 1$  for fermions. The von Neumann entropy for the mode is

$$s_{\omega,l} = [n_{\omega,l} + (-1)^\epsilon] \ln [1 + (-1)^\epsilon n_{\omega,l}] - n_{\omega,l} \ln n_{\omega,l}. \quad (43)$$

Moreover, the entropy per mode  $s_{\omega,l}$  can be put in the form of

$$s_{\omega,l} = s\left(\frac{\omega}{T_{eff}}\right), \quad (44)$$

where the  $s(x)$  is given by

$$s(x) = \frac{(-1)^\epsilon \exp x}{\exp x - (-1)^\epsilon} \ln \left[ \frac{\exp x}{\exp x - (-1)^\epsilon} \right] + \frac{\ln [\exp x - (-1)^\epsilon]}{\exp x - (-1)^\epsilon}. \quad (45)$$

Defining  $u = \frac{\omega}{T_0}$  and expanding  $s_{\omega,l}$  to  $\mathcal{O}(m_p^{-2})$  give

$$s_{\omega,l} \approx s(u) + s'(u) u \Delta, \quad (46)$$

where one has

$$\Delta = \frac{C_1 \delta_\lambda z}{2} + \frac{C_1 \delta_\omega T_0^2}{2m_p^2} u^2. \quad (47)$$

In the brick wall model, t' Hooft' found that the number of one-particle states not exceeding  $\omega$  fixed value  $l$  is

$$n(\omega, l) = \frac{1}{2\pi\hbar} \oint p_r dr, \quad (48)$$

where the integral  $\oint p_r dr$  was calculated in the Schwarzschild-like coordinate. Nevertheless, we calculate  $p_r^\pm$  in the PG coordinate in section II. In [48, 49], the integral  $\oint p_r dr$  has been found invariant under canonical transformations. Hence, the number states  $n(\omega, l)$  given in eqn. (48) is the same in the Schwarzschild-like and PG coordinates and one does not need to re-calculate it in different coordinates. Define the radial wave number  $k(r, l, \omega)$  by

$$k^\pm(r, l, \omega) = p_r^\pm, \quad (49)$$

as long as  $p_r^{\pm 2} \geq 0$ , and  $k^\pm(r, l, \omega) = 0$  otherwise. With these two Dirichlet boundaries, one finds that the number of one-particle states not exceeding  $\omega$  fixed value  $l$  is

$$n(\omega, l) = \frac{1}{2\pi\hbar} \left[ \int_{r_h+r_\varepsilon}^L k^+(r, l, \omega) dr + \int_L^{r_h+r_\varepsilon} k^-(r, l, \omega) dr \right]. \quad (50)$$

The  $p_r^\pm$  in eqn. (49) are given by eqn. (21) in section II. For a massless and neutral scalar field, one thus has

$$p_r^\pm = \frac{\pm\sqrt{\Lambda} + \sqrt{1-f}}{f} \omega \mp \frac{C_1}{m_p^2} \frac{[1 \pm \sqrt{\Lambda(1-f)}]^4 \omega^3}{f^4 \sqrt{\Lambda}} + \mathcal{O}(m_p^{-4}), \quad (51)$$

where  $\Lambda = 1 - f \frac{(l+\frac{1}{2})^2 \hbar^2}{C(r^2)\omega^2}$ .

Integrating by parts, one finds the entropy becomes

$$S = -\frac{1}{\pi\hbar} \int (2l+1) dl \int du \frac{\partial s_{\omega,l}}{\partial u} \int_{r_h+r_\varepsilon}^L k(r, l, \omega) dr, \quad (52)$$

where we define  $k(r, l, \omega) = \frac{k^+(r, l, \omega) - k^-(r, l, \omega)}{2}$ . Plugging eqns. (51) and (46) into eqn. (52) and performing the  $l$  integral which runs over the region where  $\Lambda > 0$ , we find that the entropy to  $\mathcal{O}(m_p^{-2})$  becomes

$$S \approx \frac{C(r_h^2) T_0^3}{\pi\hbar^3} \int u^2 du \int_{r_h+r_\varepsilon}^L \frac{C(r^2)}{C(r_h^2)} \frac{dr}{f^2} \left\{ 2s(u) + \frac{C_1 \delta_\omega T_0^2}{m_p^2} u^3 s'(u) + \frac{2C_1 \delta_\lambda C(r^2) T_0^2}{3f C(r_h^2) m_p^2} u^3 s'(u) - \frac{5C_1 T_0^2 u^2 s(u)}{2m_p^2 f^3} \left( 12 - \frac{48f}{5} + \frac{f^2}{2} - \frac{7f^4}{32} \right) \right\}, \quad (53)$$

where second and third terms in the bracket come from the MDR corrections to the Hawking temperature and the fourth term from the MDR corrections to  $n(\omega, l)$ . Focusing on the divergent parts near horizon, we obtain for the nonnegative integers  $a$  and  $n$

$$\int_{r_h+r_\varepsilon} \frac{C^a(r^2)}{C^a(r_h^2) f(r)^{n+1}} dr \sim \frac{1}{2^{n+1} \kappa} \sum_{k=0}^{n-1} \frac{\tilde{f}_k^{n,a}}{n-k} (\kappa r_\varepsilon)^{k-n} - \frac{\tilde{f}_n^{n,a}}{2^{n+1} \kappa} \ln \kappa r_\varepsilon, \quad (54)$$

where we expand  $f(r)^{-n}$  and  $\frac{C^a(r^2)}{C^a(r_h^2)}$  at  $r = r_h$

$$f(r)^{-n} = 2^{-n} \kappa^{-n} (r - r_h)^{-n} \sum_{j=0}^{\infty} f_j^n \kappa^j (r - r_h)^j,$$

$$\frac{C^a(r^2)}{C^a(r_h^2)} = \sum_{i=0} c_i^a \kappa^i (r - r_h)^i, \quad (55)$$

and define  $\tilde{f}_k^{n,a} = \sum_{j=0}^k f_j^{n+1} c_{k-j}^a$ . In eqn. (54), we neglect finite terms as  $\kappa r_\varepsilon \rightarrow 0$  and terms involving  $L$ . Note that we define  $\kappa = \frac{f'(r_h)}{2}$  which is the surface gravity for the black hole and hence  $T_0 = \frac{\hbar \kappa}{2\pi}$ . Thus, the divergent part of entropy near the horizon to  $\mathcal{O}(m_p^{-2})$  is

$$S \sim \frac{C(r_h^2) \kappa^2}{16\pi^4} \int s(u) u^2 du \left( \frac{1}{\kappa r_\varepsilon} - (c_1 - 2\eta) \ln \kappa r_\varepsilon \right) \\ - \frac{C(r_h^2) \kappa^2 m_p^2 \kappa^2}{16\pi^4 4\pi^2} \int s(u) u^4 du \left\{ \frac{5C_1 \delta_\omega}{2} \left( \frac{\tilde{f}_0^{1,1}}{\kappa r_\varepsilon} - \tilde{f}_1^{1,1} \ln \kappa r_\varepsilon \right) \right. \\ + \frac{5C_1 \delta_\lambda}{6} \left[ \sum_{k=0}^1 \frac{\tilde{f}_k^{2,2}}{2-k} (\kappa r_\varepsilon)^{k-2} - \tilde{f}_2^{2,2} \ln \kappa r_\varepsilon \right] \\ + 5C_1 \left[ \frac{12}{32} \sum_{k=0}^3 \frac{\tilde{f}_k^{4,1}}{4-k} (\kappa r_\varepsilon)^{k-4} - \frac{3}{5} \sum_{k=0}^2 \frac{\tilde{f}_k^{3,1}}{3-k} (\kappa r_\varepsilon)^{k-3} + \frac{\tilde{f}_0^{1,1}}{8\kappa r_\varepsilon} \right. \\ \left. \left. + \left( \frac{7\tilde{f}_0^{0,1}}{64} - \frac{\tilde{f}_1^{1,1}}{8} + \frac{3\tilde{f}_3^{3,1}}{5} - \frac{12\tilde{f}_4^{4,1}}{32} \right) \ln \kappa r_\varepsilon \right] \right\}. \quad (56)$$

## B. 2D Black Hole in Free-fall Scenario

Consider a 2D black hole with the metric of

$$ds^2 = f(r) dt^2 - \frac{dr^2}{f(r)}, \quad (57)$$

in the Schwarzschild-like coordinate. The atmosphere entropy of a massless scalar field is

$$S = \int d\omega \frac{dn(\omega, l)}{d\omega} s_\omega, \quad (58)$$

where  $s_\omega$  is the entropy per mode. Define the radial wave number  $k(r, \omega)$  by

$$k^\pm(r, \omega) = p_r^\pm, \quad (59)$$

as long as  $p_r^{\pm 2} \geq 0$ , and  $k^\pm(r, \omega) = 0$  otherwise. The  $p_r^\pm$  in eqn. (59) are given in eqn. (30) in section II. The number of one-particle states not exceeding  $\omega$  is

$$n(\omega) = \frac{1}{2\pi\hbar} \int_{r_h+r_\varepsilon}^L (k_r^+ - k_r^-) dr. \quad (60)$$

Defining the coefficients  $\sigma_s^q$  by

$$\Delta^q = \left(\frac{m_p \kappa}{2\pi}\right)^{2q} u^{2q} \sum_{s=0}^{\infty} \sigma_s^q \left(\frac{m_p \kappa}{2\pi}\right)^{2s} u^{2s}, \quad (61)$$

one has for  $s_\omega$

$$\begin{aligned} s_\omega &= s \left( \frac{\omega}{T_{eff}} \right) \\ &= \sum_{r=0}^{\infty} \sum_{q=0}^r \frac{s^{(q)}(u) u^{q+2r} \sigma_{r-q}^q \left(\frac{m_p \kappa}{2\pi}\right)^{2r}}{q!}, \end{aligned} \quad (62)$$

where  $u = \frac{\omega}{T_0}$  and we use eqn. (37) for  $T_{eff}$ . Expanding  $(1+v)^i$  at  $r = r_h$

$$(1+v)^i = \kappa^i (r - r_h)^i \sum_{j=0}^{\infty} v_j^i \kappa^j (r - r_h)^j, \quad (63)$$

we find the divergent part of entropy near the horizon becomes

$$\begin{aligned} S &\sim -\frac{1}{4\pi^2} \ln \kappa r_\varepsilon \sum_{r=0}^{\infty} \sum_{i=0}^{\infty} \sum_{q=0}^r \frac{(2i+1) \eta_i \sigma_{r-q}^q \left(\frac{m_p \kappa}{2\pi}\right)^{2i+2r}}{q!} \int du s^{(q)}(u) u^{q+2r+2i} \\ &+ \frac{1}{4\pi^2} \sum_{l=1}^{\infty} \frac{1}{l (\kappa r_\varepsilon)^l} \left(\frac{m_p \kappa}{2\pi}\right)^{2[l/3]} \sum_{r=0}^{\infty} \sum_{i=0}^{\infty} \sum_{q=0}^r \frac{\xi_{i,l} \sigma_{r-q}^q \left(\frac{m_p \kappa}{2\pi}\right)^{2i+2r}}{q!} \int du s^{(q)}(u) u^{q+2r+2i+2[l/3]}, \end{aligned} \quad (64)$$

where  $[x]$  is the smallest integer that is not less than  $x$  and we define

$$\begin{aligned} \eta_{i>0} &= \sum_{j=0}^{i-1} C_{i,j} v_{3i-j}^{-3i+j-1} \text{ and } \eta_0 = 1, \\ \xi_{i,l} &= \sum_{j=0}^{3i+3[l/3]^+-l} (2i+2[l/3]^++1) C_{i+[l/3]^+,j} v_{3i-j-l+3[l/3]^+}^{-l}. \end{aligned} \quad (65)$$

### C. 2D Black Hole in Static Scenario

Following the conventions adopted in [39], the MDR for a massless scalar particle considered here takes the form of

$$p^2 = \tilde{F}^2(E), \quad (66)$$

where we define

$$\tilde{F}(E) = E \sum_{n=0}^{\infty} \frac{\tilde{C}_n E^{2n}}{m_p^{2n}}, \quad (67)$$

with  $\tilde{C}_0 = 1$ . For a particle with the energy-momentum vector  $p_\mu$ , the energy  $\omega$  and the norm of the momentum  $p$  of the particle measured by the static observers hovering above the 2D black hole with the metric (57) are

$$\begin{aligned} E &= \frac{p_t}{\sqrt{f(r)}}, \\ p^2 &= f(r) p_r^2. \end{aligned} \quad (68)$$

Relating  $p_\mu$  to the action  $I$  by  $p_\mu = -\partial_\mu I$  gives the deformed Hamilton-Jacobi equation

$$X^2 = \tilde{F}^2(T), \quad (69)$$

where

$$T = \frac{\omega}{\sqrt{f(r)}}, \quad X = \sqrt{f(r)} p_r, \quad (70)$$

and  $\omega = \partial_t^\mu p_\mu$  is the Killing energy with respect to  $t$ . Solving eqn. (69) for  $p_r$ , one could define the radial wave number  $k(r, \omega)$  by

$$k(r, \omega) = |p_r|, \quad (71)$$

as long as  $p_r^2 \geq 0$ , and  $k(r, \omega) = 0$  otherwise. The number of one-particle states not exceeding  $\omega$  is

$$n(\omega) = \frac{1}{\pi \hbar} \int_{r_h + r_\varepsilon}^L k_r dr. \quad (72)$$

The entropy per mode  $s_\omega$  is

$$s_\omega = s\left(\frac{\omega}{T_{eff}}\right), \quad (73)$$

where  $s(x)$  is given in eqn. (45) and  $T_{eff}$  is the effective Hawking temperature. We calculated  $T_{eff}$  in [39] and it was given by

$$T_{eff} = \frac{T_0}{1 + \Delta}, \quad (74)$$

where one has

$$\Delta = \sum_{k=1}^{\infty} \eta_0^{2k} \zeta_k^0 \frac{\omega^{2k}}{m_p^{2k}}, \quad (75)$$

and  $\eta_0^{2k}$  and  $\zeta_k^0$  are defined in [39]. Defining the coefficients  $\tilde{\sigma}_s^q$  by

$$\tilde{\Delta}^q = \left(\frac{m_p \kappa}{2\pi}\right)^{2q} u^{2q} \sum_{s=0}^{\infty} \tilde{\sigma}_s^q \left(\frac{m_p \kappa}{2\pi}\right)^{2s} u^{2s}, \quad (76)$$



we find the divergent part of entropy near the horizon becomes

$$S \sim \frac{1}{4\pi^2} \sum_{n=0}^{\infty} \sum_{r=0}^{\infty} \sum_{q=0}^r \left( \frac{m_p \kappa}{2\pi} \right)^{2n+2r} \frac{2n+1}{2^n} \frac{\tilde{\sigma}_{r-q}^q}{q!} \tilde{C}_n \\ \left( \sum_{l=1}^n \frac{\tilde{f}_{n-l}^{n,0}}{l (\kappa r_\varepsilon)^l} - \tilde{f}_n^{n,0} \ln r_\varepsilon \right) \int u^{2n+2r+q} s^{(q)}(u) du, \quad (77)$$

where  $\tilde{f}_k^{n,a}$  with  $n \geq k \geq 0$  are defined in eqn. (54).

## D. Discussion

In [39] and this paper, we have calculated the divergent part of the near horizon atmosphere entropy of a massless scalar field for a 4D spherically symmetric black hole in the static and free-fall scenarios, respectively. It appears that the divergent part in both scenarios can be presented in the form of a Laurent series with respect to  $r_\varepsilon$

$$S \sim \frac{s_1^0}{\kappa r_\varepsilon} + s_0^0 \ln \kappa r_\varepsilon + \sum_{i=1}^{\infty} \frac{T_0^{2i}}{m_p^{2i}} \left( \sum_{j=1}^{\delta_i} s_j^i (\kappa r_\varepsilon)^{-j} + s_0^i \ln \kappa r_\varepsilon \right), \quad (78)$$

where  $\delta_i = 2i + 1$  in the static scenario and  $\delta_i = 3i + 1$  in the free-fall scenario. Although we calculated the atmosphere entropy around the 4D black hole to  $\mathcal{O}(m_p^{-2})$  in the free-fall scenario, the 2D black hole result suggests that eqn. (78) might hold to all orders in this scenario. For  $s_1^0$  and  $s_0^0$  in eqn. (78), we find that

$$s_1^0 = \frac{A \kappa^2}{720\pi}, \\ s_0^0 = -\frac{C(r_h^2) \kappa^2}{180} (c_1 - 2\eta), \quad (79)$$

where  $A = 4\pi C(r_h^2)$  is the horizon area. In the non-dispersive scenario ( $m_p \rightarrow \infty$ ), the terms  $\frac{s_1^0}{\kappa r_\varepsilon}$  and  $s_0^0 \ln \kappa r_\varepsilon$  are the usual leading and subleading logarithmic divergent terms, respectively. Note that  $s_1^0$  and  $s_0^0$  have already been calculated in the non-dispersive scenario[45–47].

It seems from eqn. (78) that the near horizon divergence of the atmosphere entropy gets worse for the higher order corrections in the MDR as  $\kappa r_\varepsilon \rightarrow 0$ . However, the higher order contributions in eqn. (78) are always accompanied with the powers of the factor  $\frac{T_0^2}{m_p^2} = \left( \frac{m_p \kappa}{2\pi} \right)^2$ . Thus, one might hope that the higher order divergent problem would become less severe if  $r_\varepsilon$  somehow can be related to  $m_p$ . One way to understand the value of  $r_\varepsilon$  is

introducing the proper length for  $r_\varepsilon$  as

$$\varepsilon = \int_{r_h}^{r_h+r_\varepsilon} \sqrt{g_{rr}} dr. \quad (80)$$

The brick wall is put at  $r = r_h + r_\varepsilon$  to cut off the unknown quantum physics of gravity. In this sense, the invariant distance of the wall from the horizon  $\varepsilon$  could be given by  $\varepsilon \sim m_p$ . Thus, we could define  $\alpha$  such as  $\varepsilon = \alpha m_p$ . Indeed in the 't Hooft's original calculation, equating  $\frac{s_1^0}{\kappa r_\varepsilon}$  to the Black hole's Bekenstein-Hawking entropy  $S_{BH} = \frac{A}{4m_p^2}$  gives

$$r_\varepsilon = \frac{\kappa m_p^2}{180\pi}. \quad (81)$$

Note that eqn. (80) depends on the chosen coordinate system. In the scenario without the MDR, a natural choice is that  $\varepsilon$  is measured along a static time slice. Thus, eqn. (80) is calculated in the Schwarzschild-like coordinate[45]. Assuming  $r_\varepsilon \ll r_h$ , one finds from eqn. (80)

$$r_\varepsilon \approx \frac{\kappa \varepsilon^2}{2}. \quad (82)$$

Thus, eqn. (81) gives

$$\alpha = \sqrt{\frac{1}{90\pi}}, \quad (83)$$

where we reproduce 't Hooft's result.

In the static scenario, it is still natural to assume that  $\varepsilon$  is measured along a static time slice. If we let  $\varepsilon = \alpha m_p$  in eqn. (78), we find the atmosphere entropy around the horizon becomes

$$S \sim \frac{A}{4m_p^2} \tilde{s}_0 + 2s_0^0 \ln \kappa m_p + \text{Finite terms as } m_p \kappa \rightarrow 0, \quad (84)$$

where  $\tilde{s}_0$  was given in [39]. The leading divergent coefficient  $\tilde{s}_0$  is determined by the coefficients  $\tilde{C}_n$  in the MDR (67) and  $f_j^n$  and  $c_i^a$  which are defined in eqn. (55). For a general black hole,  $f_j^n$  and  $c_i^a$  could depend on the parameters of the black hole. However, they are pure numbers for a Schwarzschild black hole. Thus, for a Schwarzschild black hole,  $\tilde{s}_0$  does not depend on the black hole's properties and the leading divergent term in eqn. (84) scales with the horizon area  $A$ .

In the free-fall scenario, one might prefer that the proper length  $\varepsilon$  is measured on a time slice orthogonal to the free-fall world lines[51]. In this case, eqn. (80) for  $\varepsilon$  should be calculated in the PG coordinate and one then has  $\varepsilon = r_\varepsilon$ . If one has  $\varepsilon = \alpha m_p$ , the

atmosphere entropy around the horizon becomes

$$S \sim \frac{A\kappa}{720\pi\alpha m_p} + s_0^0 \ln \kappa m_p + \sum_{l=1}^{\infty} \frac{s_l}{(m_p \kappa)^l} + \text{Finite terms as } m_p \kappa \rightarrow 0, \quad (85)$$

where we define  $s_l = \sum_{i=\max\{1, l-1\}}^{\infty} \frac{s_{l+2i}^i}{(2\pi)^{2i} \alpha^{l+2i}}$ . Moreover, eqn. (85) might suggest that effects of the MDR on the atmosphere entropy is nonperturbative in this case. Alternatively, inspired by the static scenario, one could choose  $r_\varepsilon$  such that the higher order terms in eqn. (78) have the same order of divergence as  $\frac{1}{\kappa r_\varepsilon}$ . Here, we could have

$$r_\varepsilon = \alpha m_p^{\frac{2}{3}} \kappa^{-\frac{1}{3}}, \quad (86)$$

where  $\alpha$  is some constant. In this case, the atmosphere entropy becomes

$$S \sim \frac{\tilde{s}_0}{\alpha (m_p \kappa)^{\frac{2}{3}}} + \frac{2s_0^0}{3} \ln m_p \kappa + \text{Finite terms as } m_p \kappa \rightarrow 0, \quad (87)$$

where we define

$$\tilde{s}_0 = s_1^0 \left[ 1 + \sum_{i=1}^{\infty} \frac{s_{3i+1}^i}{s_1^0} \left( 2\pi \alpha^{\frac{3}{2}} \right)^{-2i} \right]. \quad (88)$$

For a Schwarzschild black hole, the terms in the square bracket in eqn. (88) do not depend on the black hole's properties and the leading divergent term in eqn. (87) scales with  $A\kappa^{\frac{4}{3}} m_p^{-\frac{2}{3}}$ . In [51], the authors calculated the black hole horizon entanglement entropy for a massless scalar field with the MDR imposed in a free-fall frame. With the sub- or super-luminal dispersion with index  $n$ , they found that the entanglement entropy scales as  $A\kappa^{1+\frac{1}{n}} m_p^{-1+\frac{1}{n}}$ .

Following the argument proposed in [52], the authors in [53] obtained modified relations between the mass of a Schwarzschild black hole and its temperature and entropy. The argument connecting a MDR and some modifications of the entropy of black holes is formulated in a scheme of analysis first introduced by Bekenstein[43]. In fact, for the MDR in eqn. (1), the modified temperature of the black hole was given by

$$T_{eff} = \frac{1}{4\pi} F \left( \frac{m_p^2}{2M} \right), \quad (89)$$

where  $M$  is the mass of the black hole. The first law of black hole thermodynamics  $dS_B = \frac{dM}{T}$  and eqn. (2) lead to the modified entropy of the black hole

$$S_B = \frac{A}{4m_p^2} + 2\pi C_1 \ln \kappa m_p + \text{Finite terms as } m_p \kappa \rightarrow 0, \quad (90)$$

where  $A = 16\pi M^2$  and  $\kappa = \frac{1}{4M}$ . It is interesting to note that the modifications of the entropy of black holes in eqn. (90) are finite as  $m_p \kappa \rightarrow 0$  except the logarithmic term. If one wants the same story for the atmosphere entropy obtained in eqn. (78), one could have  $r_\varepsilon = \alpha \kappa^{\delta-1} m_p^\delta$  for some constant  $\alpha$  where  $0 < \delta \leq \frac{2}{3}$  in the static scenario and  $0 < \delta \leq \frac{1}{2}$  in the free-fall scenario. Hence, the atmosphere entropy becomes

$$S \sim \frac{s_1^0}{\alpha (\kappa m_p)^\delta} + s_0^0 \delta \ln \kappa m_p + \text{Finite terms as } m_p \kappa \rightarrow 0, \quad (91)$$

where the leading divergent term scales with  $A \kappa^{2-\delta} m_p^{-\delta}$  for a Schwarzschild black hole. The coefficient of the logarithmic term in eqn. (90) depends on  $C_1$ , a coefficient of the MDR. However, we show that the coefficient of the subleading logarithmic term in the atmosphere entropy is irrelevant to coefficients of the MDR. It only depends on the position of the wall,  $r_\varepsilon$  and the properties of the black hole.

For a 2D black hole with the Schwarzschild-like coordinate

$$ds^2 = f(r) dt^2 - \frac{dr^2}{f(r)}, \quad (92)$$

the atmosphere entropy of a massless scalar can also be presented in the form of a Laurent series with respect to  $r_\varepsilon$

$$S \sim s_0^0 \ln \kappa r_\varepsilon + \sum_{i=1}^{\infty} \frac{T_0^{2i}}{m_p^{2i}} \left( \sum_{j=1}^{\delta_i} s_j^i (\kappa r_\varepsilon)^{-j} + s_0^i \ln \kappa r_\varepsilon \right), \quad (93)$$

where  $\delta_i = 2i$  in the static scenario and  $\delta_i = 3i$  in the free-fall scenario. From eqns. (64) and (84), one has  $s_0^0 = -\frac{1}{12}$ . In the static scenario, if we assume that the proper length  $\varepsilon$  is measured along a static time slice and  $\varepsilon = \alpha m_p$ , the atmosphere entropy of a massless scalar becomes

$$S \sim -\frac{1}{6} \ln \kappa m_p + \text{Finite terms as } \kappa m_p \rightarrow 0, \quad (94)$$

where the same leading logarithmic term was also obtained in [54] for the scenario without the MDR. In the free-fall scenario, if the proper length  $\varepsilon$  is assumed to be measured on a time slice orthogonal to the free-fall world lines and we let  $\varepsilon = \alpha m_p$ , the atmosphere entropy of a massless scalar becomes

$$S \sim -\frac{1}{12} \ln \kappa m_p + \frac{1}{4\pi^2} \sum_{n=1}^{\infty} \frac{s_n}{(\kappa m_p)^n} + \text{Finite terms as } \kappa m_p \rightarrow 0, \quad (95)$$

where  $s_n$  can be determined by eqn. (64). If we want that the modifications of the entropy of black holes in eqn. (93) are finite, we could have  $r_\varepsilon = \alpha \kappa^{\delta-1} m_p^\delta$  for some constant  $\alpha$  where  $0 < \delta \leq \frac{2}{3}$  for some constant  $\alpha$ . The entropy then becomes

$$S \sim -\frac{\delta}{12} \ln \kappa m_p + \text{Finite terms as } \kappa m_p \rightarrow 0. \quad (96)$$

#### IV. BLACK HOLE EVAPORATION

In the section, we discuss the MDR effects on the evaporations of a Schwarzschild black hole to  $\mathcal{O}\left(\frac{T_h^2}{m_p^2}\right)$  in the free-fall scenario. For simplicity, we assume that the emitted particles are massless. In [55], Page counted the number of modes per frequency interval with periodic boundary conditions in a large container around the black hole and divided it by the time it takes a particle to cross the container. He then related the expected number emitted per mode  $n_{\omega,l}$  to the average emission rate per frequency interval  $\frac{dn_{\omega,l}}{dt}$  by

$$\frac{dn_{\omega,l}}{dt} = n_{\omega,l} \frac{d\omega}{2\pi\hbar}, \quad (97)$$

for each mode and frequency interval  $(\omega, \omega + d\omega)$ . Following the same argument, we find that in the MDR case

$$\frac{dn_{\omega,l}}{dt} = n_{\omega,l} \frac{\partial\omega}{\partial p_r} \frac{dp_r}{2\pi\hbar} = n_{\omega,l} \frac{d\omega}{2\pi\hbar}, \quad (98)$$

where  $\frac{\partial\omega}{\partial p_r}$  is the radial velocity of the particle and the number of modes between the wavevector interval  $(p_r, p_r + dp_r)$  is  $\frac{dp_r}{2\pi\hbar}$ . Since each particle carries off the energy  $\omega$ , the total luminosity is obtained from  $\frac{dn_{\omega,l}}{dt}$  by multiplying by the energy  $\omega$  and summing up over all energy  $\omega$  and  $l$ ,

$$L = \sum_{l=0} (2l+1) \int \omega n_{\omega,l} \frac{d\omega}{2\pi\hbar}. \quad (99)$$

However, some of the radiation emitted by the horizon might not be able to reach the asymptotic region. Before the radiation reaches the distant observer, they must pass the curved spacetime around the black hole horizon, which plays the role of a potential barrier. This effect on  $L$  can be described by a greybody factor from the scattering coefficients of the black hole. Actually, the greybody factor is given by  $|T_l(\omega)|^2$ , where  $T_l(\omega)$  represents the transmission coefficient of the black hole barrier which in general depends on the energy  $\omega$  and angular momentum  $l$  of the particle. Taking the greybody factor into account, we

find for the total luminosity

$$L = \sum_{l=0} (2l+1) \int |T_l(\omega)|^2 \omega n_{\omega,l} \frac{d\omega}{2\pi\hbar}. \quad (100)$$

The relevant radiation usually have the energy of order  $\hbar M^{-1}$  for a black hole with the mass  $M$ , one hence needs to use the wave equations given in the appendix of [39] to compute  $|T_l(\omega)|^2$  accurately. However, solving the wave equations for  $|T_l(\omega)|^2$  could be very complicated. On the other hand, we can use the geometric optics approximation to estimate  $|T_l(\omega)|^2$ . In the geometric optics approximation, we assume  $\omega \gg M$  and high energy waves will be absorbed unless they are aimed away from the black hole. Hence  $|T_l(\omega)|^2 = 1$  for all the classically allowed energy  $\omega$  and the angular momentum  $l$ , while  $|T_i(E)|^2 = 0$  otherwise. For the usual dispersion relation, the well-known Stefan's law for black holes is obtained in this approximation.

To find the classically allowed angular momentum  $l$  with fixed value of energy  $\omega$ , we consider eqn. (6) for a massless particle in the Schwarzschild black hole with the mass  $M$ . Solving eqn. (6) for  $\lambda$  to  $\mathcal{O}(m_p^{-2})$  gives

$$\frac{\lambda}{r^2} = (\omega - vp_r)^2 - \frac{2C_1}{m_p^2} (\omega - vp_r)^4 - p_r^2, \quad (101)$$

where we have  $\lambda = (l + \frac{1}{2})^2 \hbar^2$ ,  $v(r) = -\sqrt{\frac{2M}{r}}$  and  $C_1$  is given in eqn. (1). In the geometric optics approximation,  $p_r$  is always a real number. In the non-dipsersive case ( $m_p \rightarrow \infty$ ), the maximum of the RHS of eqn. (101) is  $\frac{\omega^2}{1 - \frac{2M}{r}}$ . Thus, one has

$$\lambda \leq \frac{r^2 \omega^2}{1 - \frac{2M}{r}}, \quad (102)$$

where the RHS has a minimum at  $r_{\min} = 3M$ , which is  $27M^2\omega^2$ . If the particles overcome the angular momentum barrier and get absorbed by the black hole, one must have  $\lambda \leq 27M^2\omega^2$ . In the geometric optics approximation, the Schwarzschild black hole is just like a black sphere of radius  $R = 3^{3/2}M$  [57]. When the second term in the RHS of eqn. (101) is included, the maximum of the RHS is shifted to

$$\Omega \equiv \frac{\omega^2}{1 - \frac{2M}{r}} - \frac{2C_1\omega^2}{m_p^2} \frac{\omega^2}{(1 - \frac{2M}{r})^4}. \quad (103)$$

The minimum of  $r^2\Omega$  is then shifted to

$$\lambda_{\max} \equiv 27M^2\omega^2 \left( 1 - \frac{54C_2\omega^2}{m_p^2} \right). \quad (104)$$

Therefore, the particle must have  $\lambda \leq \lambda_{\max}$  to get absorbed by the black hole. The eqn. (100) then becomes

$$L = \int \frac{\omega d\omega}{2\pi\hbar^3} \int_0^{\lambda_{\max}} n \left( \frac{\omega(1+\Delta)}{T_0} \right) d\lambda, \quad (105)$$

where eqn. (41) for  $n_{\omega,l}$  is used. For a massless particle in the Schwarzschild black hole, eqn. (24) gives

$$\Delta = -\frac{C_1}{2m_p^2} \left( 20\omega^2 - \frac{3\lambda}{4M^2} \right). \quad (106)$$

Expanding  $n \left( \frac{\omega(1+\Delta)}{T_0} \right)$  to  $\mathcal{O}(m_p^{-2})$  and then integrating eqn. (105), we find for the emission of  $n_s$  species of massless scalars and  $n_f$  species of massless spin-1/2 fermions that the total luminosity is

$$L = \frac{9m_p^2}{40960M^2} \left[ \left( n_s + \frac{7}{4}n_f \right) - C_1 (0.73n_s + 1.41n_f) \frac{m_p^2}{M^2} + \mathcal{O} \left( \frac{m_p^3}{M^3} \right) \right]. \quad (107)$$

To make a comparison with the static scenario, the results in [39] are given below for the form of MDR (1). In the static scenario, the correction to the Hawking temperature is

$$\Delta = -\frac{C_1}{2m_p^2} \left( 4\omega^2 + \frac{\lambda}{4M^2} \right), \quad (108)$$

the maximum of the angular momentum is

$$\lambda_{\max} = 27M^2\omega^2 \left( 1 - \frac{6C_1\omega^2}{m_p^2} \right), \quad (109)$$

and the total luminosity is

$$L = \frac{9m_p^2}{40960\pi M^2} \left[ \left( n_s + \frac{7}{4}n_f \right) + C_1 (0.48n_s + 0.92n_f) \frac{m_p^2}{M^2} + \mathcal{O} \left( \frac{m_p^3}{M^3} \right) \right]. \quad (110)$$

Note that the sign in front of  $C_1$  in eqn. (107) is different from that in eqn. (110). For the sub-luminal dispersion relation with  $C_1 < 0$ , it means that the total luminosity increases due to the MDR effects in the free-fall scenario while it decreases in the static scenario. In the geometric optics approximation, the black hole can be described as a black sphere for absorbing particles. The total luminosity are determined by the radius of the black sphere  $R$  and the temperature of the black hole  $T$ . Note that one has  $R = \sqrt{\frac{\lambda_{\max}}{\omega^2}}$  and  $T_{eff} \approx T_0(1 - \Delta)$ . In the static scenarios, the MDR effects increase the radius of the black sphere while they decrease the temperature of the black hole. The competition between the increased radius and the decreased temperature determines whether the luminosity would increase or decrease. It appears from eqn.(110) that the effects of decreased temperature

win the competition. In the free-fall scenario, the MDR effects also increase the radius of the black sphere. Due to the minus sign in front of  $\frac{3\lambda}{4M^2}$  in eqn. (106), the temperature of the black hole increases for  $\lambda > \frac{80}{3}M^2\omega^2$  and decreases for  $\lambda < \frac{80}{3}M^2\omega^2$ , but more slowly than in the static scenario. As a result, the effects of increased radius win the competition and hence the luminosity increases. The opposite story happens to the super-luminal case with  $C_1 > 0$ .

## V. CONCLUSION

In this paper, we used the Hamilton-Jacobi method to calculate tunneling rates of radiations across the horizon and the effective Hawking temperatures in the dispersive models with the free-fall preferred frame. After the near horizon spectrum of radiations was obtained, the thermal entropy of radiations near the horizon and luminosity of the black hole were computed. Our main results are:

- In section II, we first derived the deformed Hamilton-Jacobi equations in the dispersive models with the free-fall preferred frame. The deformed Hamilton-Jacobi equations were then solved for  $\partial_r I$  and the imaginary part of  $I$  was obtained. The corrections to the Hawking temperature were calculated for massive and charged particles to  $\mathcal{O}(m_p^{-2})$  and neutral and massless particles with  $\lambda = 0$  to all orders, respectively. It was found that corrections were suppressed by  $m_p$ .
- In section III, we used the brick wall model to compute the thermal entropy of a massless scalar field near the horizon of a 4D spherically symmetric black hole and a 2D one. For a 4D black hole, the entropy near the horizon have been calculated to  $\mathcal{O}(m_p^{-2})$  and could be written in the form of eqn. (78). The entropy became divergent as the wall approached the horizon. Various choices of the proper distance between the wall and the horizon and the corresponding entropies have been discussed. For a 2D black hole, entropies in the static and free-fall scenarios have been calculated to all orders. The leading divergent terms were logarithmic. Nevertheless, their coefficients depended on choices of the proper distance between the wall and the horizon.
- In section IV, we calculated luminosities of a Schwarzschild black hole with the mass  $M \gg m_p$ . We used the geometric optics approximation to estimate the effects of



scattering off the background. Comparison between the static scenario and the free-fall one has been given there.

## Acknowledgments

We are grateful to Houwen Wu and Zheng Sun for useful discussions. This work is supported in part by NSFC (Grant No. 11005016, 11175039 and 11375121) and the Fundamental Research Funds for the Central Universities.

- 
- [1] S. W. Hawking, “Particle Creation by Black Holes,” *Commun. Math. Phys.* **43**, 199 (1975) [Erratum-ibid. **46**, 206 (1976)].
  - [2] W. G. Unruh, “Notes on black hole evaporation,” *Phys. Rev. D* **14**, 870 (1976).
  - [3] W. G. Unruh, “Sonic analog of black holes and the effects of high frequencies on black hole evaporation,” *Phys. Rev. D* **51**, 2827 (1995).
  - [4] R. Brout, S. Massar, R. Parentani and P. Spindel, “Hawking radiation without transPlanckian frequencies,” *Phys. Rev. D* **52**, 4559 (1995) [hep-th/9506121].
  - [5] S. Corley and T. Jacobson, “Hawking spectrum and high frequency dispersion,” *Phys. Rev. D* **54**, 1568 (1996) [hep-th/9601073].
  - [6] S. Corley, “Computing the spectrum of black hole radiation in the presence of high frequency dispersion: An Analytical approach,” *Phys. Rev. D* **57**, 6280 (1998) [hep-th/9710075].
  - [7] Y. Himemoto and T. Tanaka, “A Generalization of the model of Hawking radiation with modified high frequency dispersion relation,” *Phys. Rev. D* **61**, 064004 (2000) [gr-qc/9904076].
  - [8] H. Saida and M. a. Sakagami, “Black hole radiation with high frequency dispersion,” *Phys. Rev. D* **61**, 084023 (2000) [gr-qc/9905034].
  - [9] W. G. Unruh and R. Schutzhold, “On the universality of the Hawking effect,” *Phys. Rev. D* **71**, 024028 (2005) [gr-qc/0408009].
  - [10] J. Macher and R. Parentani, “Black/White hole radiation from dispersive theories,” *Phys. Rev. D* **79**, 124008 (2009) [arXiv:0903.2224 [hep-th]].
  - [11] A. Coutant, R. Parentani and S. Finazzi, “Black hole radiation with short distance dispersion, an analytical S-matrix approach,” *Phys. Rev. D* **85**, 024021 (2012) [arXiv:1108.1821 [hep-th]].

- [12] A. Coutant and R. Parentani, “Hawking radiation with dispersion: The broadened horizon paradigm,” *Phys. Rev. D* **90**, no. 12, 121501 (2014) [arXiv:1402.2514 [gr-qc]].
- [13] F. Belgiorno, S. L. Cacciatori and F. D. Piazza, “Tunneling approach and thermality in dispersive models of analogue gravity,” arXiv:1411.7871 [gr-qc].
- [14] C. Barcelo, S. Liberati and M. Visser, “Analogue gravity,” *Living Rev. Rel.* **8**, 12 (2005) [*Living Rev. Rel.* **14**, 3 (2011)] [gr-qc/0505065].
- [15] T. Jacobson, “Trans Planckian redshifts and the substance of the space-time river,” *Prog. Theor. Phys. Suppl.* **136**, 1 (1999) [hep-th/0001085].
- [16] P. Kraus and F. Wilczek, “Selfinteraction correction to black hole radiance,” *Nucl. Phys. B* **433**, 403 (1995) [gr-qc/9408003].
- [17] P. Kraus and F. Wilczek, “Effect of selfinteraction on charged black hole radiance,” *Nucl. Phys. B* **437**, 231 (1995) [hep-th/9411219].
- [18] K. Srinivasan and T. Padmanabhan, “Particle production and complex path analysis,” *Phys. Rev. D* **60**, 024007 (1999) [gr-qc/9812028].
- [19] M. Angheben, M. Nadalini, L. Vanzo and S. Zerbini, “Hawking radiation as tunneling for extremal and rotating black holes,” *JHEP* **0505**, 014 (2005) [hep-th/0503081].
- [20] R. Kerner and R. B. Mann, “Tunnelling, temperature and Taub-NUT black holes,” *Phys. Rev. D* **73**, 104010 (2006) [gr-qc/0603019].
- [21] S. Hemming and E. Keski-Vakkuri, “The Spectrum of strings on BTZ black holes and spectral flow in the  $SL(2, R)$  WZW model,” *Nucl. Phys. B* **626**, 363 (2002) [hep-th/0110252].
- [22] A. J. M. Medved, “Radiation via tunneling from a de Sitter cosmological horizon,” *Phys. Rev. D* **66**, 124009 (2002) [hep-th/0207247].
- [23] E. C. Vagenas, “Semiclassical corrections to the Bekenstein-Hawking entropy of the BTZ black hole via selfgravitation,” *Phys. Lett. B* **533**, 302 (2002) [hep-th/0109108].
- [24] M. Arzano, A. J. M. Medved and E. C. Vagenas, “Hawking radiation as tunneling through the quantum horizon,” *JHEP* **0509**, 037 (2005) [hep-th/0505266].
- [25] S. Q. Wu and Q. Q. Jiang, “Remarks on Hawking radiation as tunneling from the BTZ black holes,” *JHEP* **0603**, 079 (2006) [hep-th/0602033].
- [26] M. Nadalini, L. Vanzo and S. Zerbini, “Hawking radiation as tunneling: The D dimensional rotating case,” *J. Phys. A* **39**, 6601 (2006) [hep-th/0511250].
- [27] B. Chatterjee, A. Ghosh and P. Mitra, “Tunnelling from black holes in the Hamilton Jacobi

- approach,” *Phys. Lett. B* **661**, 307 (2008) [arXiv:0704.1746 [hep-th]].
- [28] V. Akhmedova, T. Pilling, A. de Gill and D. Singleton, “Temporal contribution to gravitational WKB-like calculations,” *Phys. Lett. B* **666**, 269 (2008) [arXiv:0804.2289 [hep-th]].
  - [29] E. T. Akhmedov, T. Pilling and D. Singleton, “Subtleties in the quasi-classical calculation of Hawking radiation,” *Int. J. Mod. Phys. D* **17**, 2453 (2008) [arXiv:0805.2653 [gr-qc]].
  - [30] V. Akhmedova, T. Pilling, A. de Gill and D. Singleton, “Comments on anomaly versus WKB/tunneling methods for calculating Unruh radiation,” *Phys. Lett. B* **673**, 227 (2009) [arXiv:0808.3413 [hep-th]].
  - [31] R. Banerjee and B. R. Majhi, “Quantum Tunneling and Back Reaction,” *Phys. Lett. B* **662**, 62 (2008) [arXiv:0801.0200 [hep-th]].
  - [32] D. Singleton, E. C. Vagenas, T. Zhu and J. R. Ren, “Insights and possible resolution to the information loss paradox via the tunneling picture,” *JHEP* **1008**, 089 (2010) [Erratum-ibid. **1101**, 021 (2011)] [arXiv:1005.3778 [gr-qc]].
  - [33] D. Chen, H. Wu and H. Yang, “Fermion’s tunnelling with effects of quantum gravity,” *Adv. High Energy Phys.* **2013**, 432412 (2013) [arXiv:1305.7104 [gr-qc]].
  - [34] D. Chen, H. Wu and H. Yang, “Observing remnants by fermions’ tunneling,” *JCAP* **1403**, 036 (2014) [arXiv:1307.0172 [gr-qc]].
  - [35] D. Y. Chen, Q. Q. Jiang, P. Wang and H. Yang, “Remnants, fermions’ tunnelling and effects of quantum gravity,” *JHEP* **1311**, 176 (2013) [arXiv:1312.3781 [hep-th]].
  - [36] D. Chen and Z. Li, “Remarks on Remnants by Fermions’ Tunnelling from Black Strings,” *Adv. High Energy Phys.* **2014**, 620157 (2014) [arXiv:1404.6375 [hep-th]].
  - [37] D. Chen, H. Wu, H. Yang and S. Yang, *Int. J. Mod. Phys. A* **29**, no. 26, 1430054 (2014) [arXiv:1410.5071 [gr-qc]].
  - [38] B. Mu, P. Wang and H. Yang, “Minimal Length Effects on Tunnelling from Spherically Symmetric Black Holes,” *Adv. High Energy Phys.* **2015**, 898916 (2015) [arXiv:1501.06025 [gr-qc]].
  - [39] P. Wang and H. Yang, “Black Hole Radiation with Modified Dispersion Relation in Tunneling Paradigm: Static Frame,” arXiv:1505.03045 [gr-qc].
  - [40] S. Corley and T. Jacobson, “Lattice black holes,” *Phys. Rev. D* **57**, 6269 (1998) [hep-th/9709166].
  - [41] K. Srinivasan and T. Padmanabhan, “Particle production and complex path analysis,” *Phys. Rev. D* **60**, 024007 (1999) [gr-qc/9812028].

- [42] J. D. Bekenstein, “Black holes and the second law,” *Lett. Nuovo Cim.* **4**, 737 (1972).
- [43] J. D. Bekenstein, “Black holes and entropy,” *Phys. Rev. D* **7**, 2333 (1973).
- [44] S. W. Hawking, “Black Holes and Thermodynamics,” *Phys. Rev. D* **13**, 191 (1976).
- [45] G. ’t Hooft, “On the Quantum Structure of a Black Hole,” *Nucl. Phys. B* **256**, 727 (1985).
- [46] S. N. Solodukhin, “The Conical singularity and quantum corrections to entropy of black hole,” *Phys. Rev. D* **51**, 609 (1995) [hep-th/9407001].
- [47] S. N. Solodukhin, “Entanglement entropy of black holes,” *Living Rev. Rel.* **14**, 8 (2011) [arXiv:1104.3712 [hep-th]].
- [48] B. D. Chowdhury, “Problems with Tunneling of Thin Shells from Black Holes,” *Pramana* **70**, 593 (2008) [*Pramana* **70**, 3 (2008)] [hep-th/0605197].
- [49] E. T. Akhmedov, T. Pilling and D. Singleton, “Subtleties in the quasi-classical calculation of Hawking radiation,” *Int. J. Mod. Phys. D* **17**, 2453 (2008) [arXiv:0805.2653 [gr-qc]].
- [50] J. B. Hartle and S. W. Hawking, “Path Integral Derivation of Black Hole Radiance,” *Phys. Rev. D* **13**, 2188 (1976).
- [51] T. Jacobson and R. Parentani, “Black hole entanglement entropy regularized in a freely falling frame,” *Phys. Rev. D* **76**, 024006 (2007) [hep-th/0703233].
- [52] G. Amelino-Camelia, M. Arzano and A. Procaccini, “Severe constraints on loop-quantum-gravity energy-momentum dispersion relation from black-hole area-entropy law,” *Phys. Rev. D* **70**, 107501 (2004) [gr-qc/0405084].
- [53] G. Amelino-Camelia, M. Arzano, Y. Ling and G. Mandanici, “Black-hole thermodynamics with modified dispersion relations and generalized uncertainty principles,” *Class. Quant. Grav.* **23**, 2585 (2006) [gr-qc/0506110].
- [54] R. B. Mann, L. Tarasov and A. Zelnikov, “Brick walls for black holes,” *Class. Quant. Grav.* **9**, 1487 (1992).
- [55] D. N. Page, “Particle Emission Rates from a Black Hole: Massless Particles from an Uncharged, Nonrotating Hole,” *Phys. Rev. D* **13**, 198 (1976).
- [56] S. Corley and T. Jacobson, “Hawking spectrum and high frequency dispersion,” *Phys. Rev. D* **54**, 1568 (1996) [hep-th/9601073].
- [57] R. M. Wald, “General Relativity,” Chicago, Usa: Univ. Pr. ( 1984) 491p